

# RaspGel-B

## Oral Anesthetic

Benzocaine Gel USP 20%

Storage : 20°C - 25°C (68°F - 77°F)

**COMPOSITION:** Benzocaine USP 20% w/w in a water soluble base consisting of polyethylene glycol 300, polyethylene glycol 3350, sodium saccharin, Raspberry flavour and Ponceau 4R.

For professional use only in the practice of dentistry  
Do not use if protective seal under cap is broken or damaged.

**INDICATIONS:** For temporary pain relief of oral mucosa due to minor irritation or injury of the mouth and gums or of pain associated with canker sores, dentures or orthodontic appliances. For topical application to oral mucosa only.

**SPECIAL PRECAUTIONS DURING HANDLING AND STORAGE:** PATIENT SHOULD NOT INGEST OR SWALLOW. AVOID CONTACT WITH EYES. If this happens, rinse thoroughly with water. Hypersensitivity/allergy has been known to occur, in which case, discontinue use. Localized allergic reactions may occur after prolonged or repeated use. Do not use on patients with history of allergy due to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics. Discontinue use if the following symptoms appear: weakness, confusion, headache, difficulty in breathing and/or pale, gray or blue coloured skin as these may be signs of methemoglobinemia, a rare disorder, which may appear up to 2 hours after use. Use smallest amount possible to achieve desired result. For occasional use only. Use with caution in children.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Centre right away.

Apply to the affected area, wait until relief is obtained, and rinse the mouth before reinserting the appliance.

Contact a dentist at regular intervals when using this product to relieve pain during adjustment of new dentures or other dental appliances.

Do not eat for one hour following use.

Do not chew gum or food while numbness persists.

Store in original container protected from direct sunlight.

Keep container tightly closed and sealed until ready for use.

Containers that have been opened must be carefully resealed and kept upright to prevent leakage.

Avoid prolonged periods of storage at higher temperatures, as it may cause undesirable degradation to products.

### PRESENTATIONS

Benzocaine Gel USP 20% is available in 30 g jar

Manufactured and Marketed by:

**SEPTODONT HEALTHCARE INDIA PVT. LTD.**

C-1/2, MIDC, Taloja, Tai. Panvel,

Dist: Raigad - 410 208,

Maharashtra (India)

Customer Care No.: +91 8657774545

info@septodontindia.com



R142112250